

PRIOR AUTHORIZATION POLICY

POLICY: Homozygous Familial Hypercholesterolemia – Evkeeza Prior Authorization Policy

- Evkeeza[®] (evinacumab-dgnb intravenous infusion – Regeneron)

REVIEW DATE: 05/08/2024

OVERVIEW

Evkeeza, an angiopoietin-like 3 inhibitor, is indicated as an adjunct to other low-density lipoprotein cholesterol (LDL-C) lowering therapies for the treatment of **homozygous familial hypercholesterolemia** (HoFH) in patients ≥ 5 years of age.¹

In the pivotal trial that led to approval of Evkeeza, patients were receiving additional medications to lower LDL-C levels such as statins (94% [77% of patients at high-intensity statin doses]), a proprotein convertase subtilisin kexin type 9 (PCSK9) inhibitor (77%), ezetimibe (75%), and Juxtapid[®] (lomitapide capsules). Although some Phase II data are available,³ the safety and effectiveness of Evkeeza have not been established in patients with other causes of hypercholesterolemia, including those with heterozygous familial hypercholesterolemia (HeFH).¹ The effects of Evkeeza on cardiovascular (CV) morbidity and mortality have not been determined.

Disease Overview

Familial hypercholesterolemias, which include HeFH and HoFH, encompass a group of genetic defects that cause severe elevations in LDL-C levels, as well as other lipid parameters.^{4,5} HoFH impacts approximately 1 in 300,000 to 1,000,000 persons. The condition is most commonly due to impaired functionality of the low-density lipoprotein (LDL) receptor which leads to a low or absence of clearance of LDL-C from the circulation. Currently known causes of familial hypercholesterolemia include mutations in the LDL receptor, apolipoprotein B, or PCSK9 genes. Patients with familial hypercholesterolemia may have physical findings such as tendon or cutaneous xanthomas, which may occur in childhood. Individuals with familial hypercholesterolemia are at very high risk of atherosclerotic cardiovascular disease (ASCVD) at a premature age. Most guidelines recommended LDL-C targets of < 100 mg/dL for adults and < 70 mg/dL for adults with ASCVD or other risk factors. Statins are the initial treatment for familial hypercholesterolemia. For patients with HoFH, high-intensity statin therapy is recommended, with ezetimibe added as well. Therapy with a PCSK9 inhibitor (e.g., Repatha[®] [evolocumab subcutaneous injection], Praluent[®] [alirocumab subcutaneous injection]) is usually the next step. Juxtapid can be added onto maximal lipid-lowering therapy and Evkeeza may be considered. Combination therapy is required for most patients. LDL apheresis is recommended in certain circumstances. The diagnosis of HoFH can be done by genetic or clinical criteria.⁵ An untreated LDL-C (> 400 mg/dL) is suggestive of HoFH. Patients may have cutaneous or tendon xanthomas before 10 years of age and/or untreated elevated LDL-C levels consistent with HeFH in both parents. In the digenic form, one parent may have normal LDL-C levels and the other may have LDL-C levels consistent with HoFH.

Guidelines

Guidelines provide strategies for managing familial hypercholesterolemia, including HoFH, and mention the role of Evkeeza.^{5,6}

- **American College of Cardiology (2022):** Specialized therapies, one of which includes Evkeeza, may be needed to control LDL-C in certain patients (e.g., those with HoFH) who have had an inadequate response to statins, with or without ezetimibe, and PCSK9 inhibitors.⁶

05/08/2024

© 2024. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

- **European Atherosclerosis Society (2023):** Clinical guidance by this organization recommends lipid-lowering therapy be initiated with high-intensity statin therapy and ezetimibe.⁵ A PCSK9 inhibitor can be added as well. If not at LDL-C goals, other agents can be alternatives as well (e.g., Juxtapid, Evkeeza). Lipoprotein apheresis may also be considered. The goal is to reduce LDL-C to < 115 mg/dL in children and adolescents, < 70 mg/dL in adults if no major ASCVD risk factors are present, and < 55 mg/dL if patients have ASCVD or major ASCVD risk factors.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Evkeeza. All approvals are provided for the duration noted below. A patient who has previously met Initial Therapy criteria for Evkeeza for the requested indication under the Coverage Review Department and is currently receiving Evkeeza is only required to meet continuation of therapy criteria (i.e., currently receiving therapy). If past criteria has not been met under the Coverage Review Department and the patient is currently receiving Evkeeza, or is restarting Evkeeza, Initial Therapy criteria must be met.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Evkeeza is recommended in those who meet the following criteria:

FDA-Approved Indication

1. **Homozygous Familial Hypercholesterolemia.** Approve for 1 year if the patient meets ONE of the following (A or B):

A) **Initial Therapy.** Approve if the patient meets ALL of the following (i, ii, iii, and iv):

i. Patient is ≥ 5 years of age; AND

ii. Patient meets ONE of the following (a, b, or c):

a) Patient has phenotypic confirmation of homozygous familial hypercholesterolemia; OR
Note: Examples include pathogenic variants at the low-density lipoprotein receptor (LDLR), apolipoprotein B (apo B), proprotein convertase subtilisin kexin type 9 (PCSK9), or low-density lipoprotein receptor adaptor protein 1 (LDLRAP1) gene.

b) Patient has an untreated low-density lipoprotein cholesterol (LDL-C) level > 400 mg/dL AND meets ONE of the following [(1) or (2)]:

Note: Untreated refers to prior to therapy with any antihyperlipidemic agent.

(1) Patient had clinical manifestations of homozygous familial hypercholesterolemia before the age of 10 years; OR

Note: Clinical manifestations of homozygous familial hypercholesterolemia are cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas, or xanthelasma.

(2) At least one parent of the patient had untreated LDL-C levels or total cholesterol levels consistent with familial hypercholesterolemia; OR

Note: An example of familial hypercholesterolemia is an untreated LDL-C level ≥ 190 mg/dL and/or an untreated total cholesterol level > 250 mg/dL.

c) Patient has a treated LDL-C level ≥ 300 mg/dL AND meets ONE of the following [(1) or (2)]:

Note: Treated refers to after therapy with at least one antihyperlipidemic agent. Some examples of antihyperlipidemic agents include statins (e.g., atorvastatin, rosuvastatin, lovastatin, simvastatin, pravastatin), ezetimibe, a PCSK9 inhibitor (i.e., Repatha

[evolocumab subcutaneous injection, Praluent [alirocumab subcutaneous injection]], or Juxtapid (lomitapide capsules).

- (1) Patient had clinical manifestations of homozygous familial hypercholesterolemia before the age of 10 years; OR

Note: Examples of clinical manifestations of homozygous familial hypercholesterolemia are cutaneous xanthomas, tendon xanthomas, arcus cornea, tuberous xanthomas, or xanthelasma.

- (2) At least one parent of the patient had untreated LDL-C levels or total cholesterol levels consistent with familial hypercholesterolemia; AND

Note: An example of familial hypercholesterolemia is an untreated LDL-C ≥ 190 mg/dL and/or an untreated total cholesterol > 250 mg/dL.

iii. Patient meets ONE of the following (a or b):

- a) Patient meets ALL of the following [(1), (2), and (3)]:

- (1) Patient has tried one high-intensity statin therapy (i.e., atorvastatin ≥ 40 mg daily; rosuvastatin ≥ 20 mg daily [as a single entity or as a combination product]); AND

- (2) Patient has tried one high-intensity statin along with ezetimibe (as a single-entity or as a combination product) for ≥ 8 continuous weeks; AND

- (3) Low-density lipoprotein cholesterol level after this treatment regimen remains ≥ 70 mg/dL; OR

- b) Patient has been determined to be statin intolerant by meeting ONE of the following [(1) or (2)]:

- (1) Patient experienced statin-related rhabdomyolysis; OR

Note: Rhabdomyolysis is statin-induced muscle breakdown that is associated with markedly elevated creatine kinase levels (at least 10 times the upper limit of normal), along with evidence of end organ damage which can include signs of acute renal injury (noted by substantial increases in serum creatinine [Scr] levels [a ≥ 0.5 mg/dL increase in Scr or doubling of the Scr] and/or myoglobinuria [myoglobin present in urine]).

- (2) Patient meets ALL of the following [(a), (b), and (c)]:

- (a) Patient experienced skeletal-related muscle symptoms; AND

Note: Examples of skeletal-related muscle symptoms include myopathy (muscle weakness) or myalgia (muscle aches, soreness, stiffness, or tenderness).

- (b) The skeletal-muscle related symptoms occurred while receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or combination products); AND

- (c) When receiving separate trials of both atorvastatin and rosuvastatin (as single-entity or as combination products) the skeletal-related muscle symptoms resolved upon discontinuation of each respective statin therapy (atorvastatin and rosuvastatin); AND

Note: Examples of skeletal-related muscle symptoms include myopathy and myalgia.

iv. Patient meets ONE of the following (a, b, or c):

- a) Patient meets BOTH of the following [(1) and (2)]:

- (1) Patient has tried one PCSK9 inhibitor for ≥ 8 continuous weeks; AND

Note: Examples of PCSK9 inhibitors include Repatha (evolocumab subcutaneous injection) and Praluent (alirocumab subcutaneous injection).

- (2) The LDL-C level after this PCSK9 inhibitor therapy remains ≥ 70 mg/dL; OR

- b) Patient is known to have two LDL-receptor negative alleles; OR

- c) Patient is 5 to 9 years of age; OR

B) Patient Currently Receiving Evkeeza. Approve if according to the prescriber, the patient has experienced a response to therapy.

Note: Examples of a response to therapy include decreasing LDL-C, total cholesterol, non-high-density lipoprotein (non-HDL-C), or apolipoprotein B levels. Also, if the patient is currently

receiving the requested therapy but has not previously received approval of Evkeeza for this specific indication through the Coverage Review Department, review under criteria for Initial Therapy. If the patient is restarting therapy with Evkeeza, Initial Therapy criteria must be met.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Evkeeza is not recommended in the following situations:

1. **Heterozygous Familial Hypercholesterolemia.** The safety and effectiveness of Evkeeza have not been established in patients with hypercholesterolemia who do not have HoFH, including those with HeFH.¹
2. **Hyperlipidemia.** Although data are available, the prescribing information for Evkeeza states that the safety and efficacy of Evkeeza have not been established in patients with other forms of hypercholesterolemia.^{1,3}
Note: This is not associated with HoFH and is referred to as combined hyperlipidemia, hypercholesterolemia (pure, primary), dyslipidemia, or increased/elevated LDL-C levels.
3. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

1. Evkeeza® intravenous infusion [prescribing information]. Tarrytown, NY: Regeneron; March 2023.
2. Raal FJ, Rosenson RS, Reeskamp LF, et al, for the ELIPSE HoFH investigators. Evkeeza for homozygous familial hypercholesterolemia. *N Engl J Med.* 2020;383(8):711-720.
3. Rosenson RS, Burgess LJ, Ebenbichler CF, et al. Evkeeza in patients with refractory hypercholesterolemia. *N Engl J Med.* 2020;383(24):2307-2319.
4. Raal FJ, Hovingh GK, Catapano AL. Familial hypercholesterolemia treatments: guidelines and new therapies. *Atherosclerosis.* 2018;277:483-492.
5. Cuchel M, Raal FJ, Hegele RA, et al. 2023 update on European Atherosclerosis Society Consensus Statement on Homozygous Familial Hypercholesterolaemia: new treatments and clinical guidance. *Eur Heart J.* 2023;44:2277-2291.
6. Lloyd-Jones DM, Morris PB, Ballantyne CM, et al. 2022 ACC Expert Consensus Decision Pathway on the Role of Non-Statins Therapies for LDL-Cholesterol Lowering in the Management of Atherosclerotic Cardiovascular Disease Risk. *J Am Coll.* 2022;80(14):1366-1418.

