

PRIOR AUTHORIZATION POLICY

POLICY: Oncology (Injectable) – Monjuvi Prior Authorization Policy

- Monjuvi® (tafasitamab-cxix intravenous infusion – MorphoSys/Incyte)

REVIEW DATE: 09/04/2024

OVERVIEW

Monjuvi, a CD19-directed antibody-drug conjugate, is indicated in combination with lenalidomide for the treatment of relapsed or refractory **diffuse large B-cell lymphoma** (DLBCL) not otherwise specified, including DLBCL arising from low grade lymphoma, and who are not eligible for autologous stem cell transplant in adults.¹

Dosing Information

Monjuvi is administered as a weight-based intravenous infusion.¹ It should be given in combination with lenalidomide for a maximum of 12 cycles, then as monotherapy until disease progression or unacceptable toxicity.

Guidelines

The National Comprehensive Cancer Network (NCCN) guidelines for B-cell lymphomas (version 3.2024 – August 26, 2024) include Monjuvi + lenalidomide among the alternatives for second-line and subsequent therapy of DLBCL, histologic transformation of indolent lymphomas to DLBCL, human immunodeficiency virus (HIV)-related B-cell lymphoma, post-transplant lymphoproliferative disorders, and high-grade B-cell lymphoma.^{2,3} NCCN also notes that it is unclear if Monjuvi would have a negative impact on the efficacy of subsequent anti-CD19 CAR T-cell therapy.

POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Monjuvi. All approvals are provided for the duration noted below. Because of the specialized skills required for evaluation and diagnosis of patients treated with Monjuvi as well as the monitoring required for adverse events and long-term efficacy, approval requires Monjuvi to be prescribed by or in consultation with a physician who specializes in the condition being treated.

Automation: None.

RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Monjuvi is recommended in those who meet one of the following criteria:

FDA-Approved Indication

- 1. Diffuse Large B-Cell Lymphoma.** Approve for 1 year if the patient meets ALL of the following (A, B, C, D, and E):
 - A) Patient is ≥ 18 years of age; AND
 - B) Patient has been treated with at least one prior chemotherapy regimen; AND
 - C) According to the prescriber, the patient is not eligible for autologous stem cell transplant; AND
 - D) Patient meets ONE of the following (i or ii):

09/04/2024

© 2024. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

- i. Monjuvi will be used in combination with Revlimid (lenalidomide capsules); OR
 - ii. Patient has already received 12 cycles of Monjuvi; AND
- E) The medication is prescribed by or in consultation with an oncologist.

Other Uses with Supportive Evidence

2. **B-Cell Lymphoma.** Approve for 1 year if the patient meets ALL of the following (A, B, C, D, and E):
Note: Examples include high-grade B-cell lymphoma, HIV-related B-cell lymphoma, post-transplant lymphoproliferative disorders, histologic transformation of indolent lymphomas to diffuse large B-cell lymphoma.
- A) Patient is ≥ 18 years of age; AND
 - B) Patient has been treated with at least one prior chemotherapy regimen; AND
Note: Examples of chemotherapy regimens include CHOP (cyclophosphamide, doxorubicin, vincristine, prednisone) plus rituximab or Gazyva (obinutuzumab intravenous infusion), CVP (cyclophosphamide, vincristine, prednisone) plus rituximab or Gazyva, or lenalidomide plus rituximab.
 - C) According to the prescriber, the patient is not eligible for autologous stem cell transplant; AND
 - D) Patient meets ONE of the following (i or ii):
 - i. Monjuvi will be used in combination with Revlimid (lenalidomide capsules); OR
 - ii. Patient has already received 12 cycles of Monjuvi; AND
 - E) The medication is prescribed by or in consultation with an oncologist.

CONDITIONS NOT RECOMMENDED FOR APPROVAL

Coverage of Monjuvi is not recommended in the following situations:

- 1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

REFERENCES

- 1. Monjuvi® intravenous infusion [prescribing information]. Boston, MA: MorphoSys/Incyte; May 2024.
- 2. The NCCN B-Cell Lymphoma Clinical Practice Guidelines in Oncology (version 3.2024 – August 26, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on August 30, 2024.
- 3. The NCCN Drugs & Biologics Compendium. © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on August 30, 2024. Search term: tafasitamab.