

## PRIOR AUTHORIZATION POLICY

**POLICY:** Oncology (Injectable) – Mylotarg Prior Authorization Policy

- Mylotarg<sup>™</sup> (gemtuzumab ozogamicin intravenous infusion – Pfizer)

**REVIEW DATE:** 07/17/2024

---

### OVERVIEW

Mylotarg, an antibody-drug conjugate directed towards the CD33 antigen, is indicated for the following:<sup>1</sup>

- **CD33-positive acute myeloid leukemia (AML)**, newly diagnosed, in adults and pediatric patients  $\geq 1$  month of age; AND
- **CD33-positive AML**, relapsed or refractory, in adults and in pediatric patients  $\geq 2$  years of age.

### Guidelines

The National Comprehensive Cancer Network (NCCN) guidelines for **AML** (version 3.2024 – May 17, 2024) recommend Mylotarg for induction therapy, post-remission therapy, and for relapsed/refractory CD33-positive AML.<sup>2,3</sup> Mylotarg can be used as a single agent or in combination with cytarabine and daunorubicin. The NCCN guidelines for AML also recommend Mylotarg in patients  $\geq 18$  years of age for induction and consolidation therapy for acute promyelocytic leukemia, and for relapsed disease. Mylotarg can be used in combination with tretinoin and/or arsenic trioxide.

### POLICY STATEMENT

Prior Authorization is recommended for prescription benefit coverage of Mylotarg. All approvals are provided for the duration noted below. In cases where the approval is authorized in months, 1 month is equal to 30 days. Because of the specialized skills required for evaluation and diagnosis of patients treated with Mylotarg as well as the monitoring required for adverse events and long-term efficacy, approval requires Mylotarg to be prescribed by or in consultation with a physician who specializes in the condition being treated.

**Automation:** None.

### RECOMMENDED AUTHORIZATION CRITERIA

Coverage of Mylotarg is recommended in those who meet one of the following criteria:

#### FDA-Approved Indication

1. **Acute Myeloid Leukemia.** Approve for the duration noted if the patient meets ONE of the following (A or B):
  - A) Newly diagnosed CD33-positive disease: Approve for 1 year if the patient meets BOTH of the following (i and ii):
    - i. Patient is  $\geq 1$  month of age; AND
    - ii. Mylotarg is prescribed by or in consultation with an oncologist; OR
  - B) Relapsed or refractory CD33-positive disease: Approve for 1 month if the patient meets BOTH of the following (i and ii):
    - i. Patient is  $\geq 2$  years of age; AND
    - ii. Mylotarg is prescribed by or in consultation with an oncologist.

07/17/2024

© 2024. All Rights Reserved.

This document is confidential and proprietary. Unauthorized use and distribution are prohibited.

### **Other Uses with Supportive Evidence**

2. **Acute Promyelocytic Leukemia.** Approve for 6 months if the patient meets BOTH of the following (A and B):
  - A) Patient is  $\geq 18$  years of age; AND
  - B) Mylotarg is prescribed by or in consultation with an oncologist.

### **CONDITIONS NOT RECOMMENDED FOR APPROVAL**

Coverage of Mylotarg is not recommended in the following situations:

1. Coverage is not recommended for circumstances not listed in the Recommended Authorization Criteria. Criteria will be updated as new published data are available.

### **REFERENCES**

1. Mylotarg™ intravenous infusion [prescribing information]. Philadelphia, PA: Pfizer; August 2021.
2. The NCCN Acute Myeloid Leukemia Clinical Practice Guidelines in Oncology (version 3.2024 – May 17, 2024). © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on July 9, 2024.
3. The NCCN Drugs and Biologics Compendium. © 2024 National Comprehensive Cancer Network. Available at: <http://www.nccn.org>. Accessed on July 9, 2024. Search term: gemtuzumab.